

1/56

FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6



09049865-032798

7/56

FIG. 7



86220"59864060

FIG. 8




FIG. 9




FIG. 10




FIG. 11




FIG. 12




FIG. 13



05/22/00 55661060

FIG. 14






FIG. 15




060498E : 0279

FIG. 18



SECRET



20/56

FIG. 20



FIG. 21




FIG. 22



FIG. 23



FIG. 24




FIG. 25A




FIG. 25B





FIG. 25C






FIG. 25D

FIG. 26



59864060

FIG. 27






FIG. 28A





FIG. 28B

FIG. 29B



B

FIG. 29A



30/56

FIG. 30




FIG. 31



000000-022-59864060

31/56

FIG. 32




FIG. 33



554220 59864050

FIG. 34 Microencapsulated Neonatal Pig Islet Transplants Into Diabetic NOD Mice
Treated with CTLA4-Ig for 21 Days




FIG. 35 Microencapsulated Neonatal Pig Islet Transplants into Diabetic NOD Mice
Treated with mutant CTLA4-Ig for 21 Days




FIG. 36

Neonatal Porcine Islet Xenografts in NOD Mice: Effects of CTLA-4Ig and Microencapsulation




FIG. 37




FIG. 38

Spontaneous Proliferation by SPC from
NODs with rejected or functioning grafts




FIG. 39A IL-2 present in peritoneal fluid on sac day
Transplanted NODs




FIG. 39B IL-2 present in peritoneal fluid on sac day
Untransplanted mice




FIG. 40A IL-2 secreted by SPC cultured with porcine islets
Transplanted NODs - Not Rejecting




FIG. 40B IL-2 secreted by SPC cultured with porcine islets
Transplanted NODs - Rejecting




FIG. 41A IFN-gamma present in peritoneal fluid on sac day
Transplanted NODs




FIG. 41B IFN-gamma present in peritoneal fluid on sac day
Untransplanted mice




FIG. 42A IFN gamma secreted by SPC cultured with porcine islets
Transplanted NODs - Not Rejecting




FIG. 42B IFN-gamma secretion by SPC cultured with porcine islets
Transplanted NODs - Rejecting




FIG. 43A IL-4 present in peritoneal fluid on sac day
Transplanted NODs




FIG. 43B IL-4 present in peritoneal fluid on sac day
Untransplanted mice




FIG. 44A IL-4 secreted by SPC cultured with porcine islets
Transplanted NODs - Not Rejecting




FIG. 44B IL-4 secreted by SPC cultured with porcine islets
Transplanted NODs - Rejecting




FIG. 45A IL-5 present in peritoneal fluid on sac day
Transplanted NODs




FIG. 45B IL-5 present in peritoneal fluid on sac day
Untransplanted mice




FIG. 46A IL-10 present in peritoneal fluid on sac day
Transplanted NODs




FIG. 46B IL-10 present in peritoneal fluid on sac day
Untransplanted mice




FIG. 47A IL-10 secreted by SPC cultured with porcine islets
Transplanted NODs - Not Rejecting




FIG. 47B IL-10 secreted by SPC cultured with porcine islets
Transplanted NODs - Rejecting




FIG. 48A IL-12 present in peritoneal fluid on sac day
Transplanted NODs




FIG. 48B IL-12 present in peritoneal fluid on sac day
Untransplanted mice




FIG. 49A IL-12 secreted by SPC cultured with porcine islets
Transplanted NODs - Not Rejecting




FIG. 49B IL-12 secreted by SPC cultured with porcine islets
Transplanted NODs - Rejecting




FIG. 50A TNF alpha present in peritoneal fluid on sac day
Transplanted NODs




FIG. 50B TNF alpha present in peritoneal fluid on sac day
Untransplanted NODs




FIG. 51A TNF- α secreted by SPC cultured with porcine islets
Transplanted NODs - Not Rejecting




FIG. 51B TNF- α secreted by SPC cultured with porcine islets
Transplanted NODs - Rejecting




FIG. 52A TGF beta present in peritoneal fluid on sac day
Transplanted NODs




FIG. 52B TGF beta present in peritoneal fluid on sac day
Untransplanted mice




FIG. 53A TGF- β secreted by SPC cultured with porcine islets
Transplanted NODs - Not Rejecting




FIG. 53B TGF- β secreted by SPC cultured with porcine islets
Transplanted NODs - Rejecting




FIG. 54A NO₂ produced by SPCs cultured with pig islets
Transplanted NODs - Not Rejecting




FIG. 54B Transplanted NODs - Rejecting




FIG. 55A

NO₂ produced by PECs after 96 hr in culture
with original encapsulated porcine islets

Transplanted NODs - Not Rejecting




FIG. 55B

Transplanted NODs - Rejecting




FIG. 56

Proliferation of SPC from NOD #1335 to pig islets or insulin
(Expt 288)




FIG. 57A Proliferation of SPC from NOD #1335
to pig islet cells in vitro (Expt 288)




FIG. 57B Proliferation of SPC from NOD #1335
Expt 288




FIG. 58 NO₂ Production by Cells from NOD #1335 (Expt 288)

